arXiv:math/9911114v1 [math.QA] 16 Nov 1999

Nonstandard q-deformation of the universal enveloping algebra $\mathbf{U(so_n)}$

A. U. Klimyk

Bogolyubov Institute for Theoretical Physics
Metrologichna str. 14-B
Kiev 03143, Ukraine

Abstract

We describe properties of the nonstandard q-deformation $U'_q(\mathrm{so}_n)$ of the universal enveloping algebra $U(\mathrm{so}_n)$ of the Lie algebra so_n which does not coincide with the Drinfeld–Jimbo quantum algebra $U_q(\mathrm{so}_n)$. Irreducible representations of this algebras for q a root of unity $q^p=1$ are given. These representations act on p^N-dimensional linear space (where N is a number of positive roots of the Lie algebra so_n) and are given by $r=\dim \mathrm{so}_n$ complex parameters.

We consider a q-deformation $U'_q(\mathrm{so}_n)$ of the universal enveloping algebra $U(\mathrm{so}_n)$ of the Lie algebra so_n which does not coincide with the Drinfeld–Jimbo quantum algebra $U_q(\mathrm{so}_n)$. The algebra $U'_q(\mathrm{so}_n)$ is constructed without using the Cartan subalgebra and roots. This algebra has no root elements.

The Drinfeld–Jimbo algebra $U_q(\mathrm{so}_n)$ is obtained by deforming Serre's relations for generating elements $E_1,\cdots,E_l,\ F_1,\cdots,F_l,\ H_1,\cdots,H_l$ of $U_q(\mathrm{so}_n)$. In order to obtain $U'_q(\mathrm{so}_n)$ we have to take determining relations for the generating elements I_{21}, $I_{32},\cdots,I_{n,n-1}$ of $U_q(\mathrm{so}_n)$ (they do not coincide with E_j, F_j, H_j) and to deform these relations. The elements $I_{21},\ I_{32},\cdots,I_{n,n-1}$ belong to the basis I_{ij}, $i>j$, of the Lie algebra so_n. The matries I_{ij}, $i>j$, are defined as $I_{ij}=E_{ij}-E_{ji}$, where E_{ij} is the matrix with entries $(E_{ij})_{rs}=\delta_{ir}\delta_{js}$. The universal enveloping algebra $U(\mathrm{so}_n)$ is generated by a part of the basis elements I_{ij}, $i>j$, namely, by the elements I_{21}, $I_{32},\cdots,I_{n,n-1}$. These elements satisfy the relations

$$I^2_{i,i-1}I_{i+1,i}-2I_{i,i-1}I_{i+1,i}I_{i,i-1}+I_{i+1,i}I^2_{i,i-1}=-I_{i+1,i},$$

$$I_{i,i-1}I^2_{i+1,i} - 2I_{i+1,i}I_{i,i-1}I_{i+1,i} + I^2_{i+1,i}I_{i,i-1} = -I_{i,i-1},$$

$$I_{i,i-1}I_{j,j-1} - I_{j,j-1}I_{i,i-1} = 0 \quad \text{for} \quad |i-j| > 1.$$

Theorem 1. *The universal enveloping algebra $U(\mathrm{so}_n)$ is isomorphic to the complex associative algebra (with a unit element) generated by the elements I_{21}, $I_{32}, \cdots, I_{n,n-1}$ satisfying the above relations.*

We make the q-deformation of these relations by $2 \to [2] := (q^2 - q^{-2})/(q - q^{-1}) = q + q^{-1}$. As a result, we obtain the complex associative algebra generated by elements I_{21}, $I_{32}, \cdots, I_{n,n-1}$ satisfying the relations

$$I^2_{i,i-1}I_{i+1,i} - (q+q^{-1})I_{i,i-1}I_{i+1,i}I_{i,i-1} + I_{i+1,i}I^2_{i,i-1} = -I_{i+1,i},$$

$$I_{i,i-1}I^2_{i+1,i} - (q+q^{-1})I_{i+1,i}I_{i,i-1}I_{i+1,i} + I^2_{i+1,i}I_{i,i-1} = -I_{i,i-1},$$

$$I_{i,i-1}I_{j,j-1} - I_{j,j-1}I_{i,i-1} = 0 \quad \text{for} \quad |i-j| > 1.$$

This algebra was introduce by us in [1] and is denoted by $U'_q(\mathrm{so}_n)$. There are the following motivations for studying this algebra and its representations:

1. The algebra of observables in 2+1 quantum gravity is isomorphic to the algebra $U'_q(\mathrm{so}_n)$ or to its quotient algebra [2, 3].

2. A quantum analogue of the Riemannian symmetric space $SU(n)/SO(n)$ is constructed by means of the algebra $U'_q(\mathrm{so}_n)$ (see [4]).

3. A q-analogue of the theory of harmonic polynomials (q-harmonic polynomials on quantum vector space $\mathbf{C}^n_q$) is constructed by using the algebra $U'_q(\mathrm{so}_n)$ (see [5]).

In $U'_q(\mathrm{so}_n)$ we can determine [6] elements analogous to the matrices I_{ij}, $i > j$, of the Lie algebra so_n. In order to give them we use the notation $I_{k,k-1} \equiv I^+_{k,k-1} \equiv I^-_{k,k-1}$. Then for $k > l+1$ we define recursively

$$I^{\pm}_{kl} := [I_{l+1,l}, I_{k,l+1}]_{q^{\pm 1}} \equiv q^{\pm 1/2} I_{l+1,l}I_{k,l+1} - q^{\mp 1/2} I_{k,l+1}I_{l+1,l}.$$

(Note that similar sets of elements of $U'_q(\mathrm{so}_n)$ are also introduced in [5]). The elements I^+_{kl}, $k > l$, satisfy the commutation relations

$$[I^+_{ln}, I^+_{kl}]_q = I^+_{kn}, \quad [I^+_{kl}, I^+_{kn}]_q = I^+_{ln}, \quad [I^+_{kn}, I^+_{ln}]_q = I^+_{kl} \quad \text{for} \quad k > l > n,$$

$$[I^+_{kl}, I^+_{nr}] = 0 \quad \text{for} \quad k > l > n > r \;\; \text{and} \;\; k > n > r > l,$$

$$[I^+_{kl}, I^+_{nr}]_q = (q - q^{-1})(I^+_{lr}I^+_{kn} - I^+_{kr}I^+_{nl}) \quad \text{for} \quad k > n > l > r.$$

For I^-_{kl}, $k > l$, the commutation relations are obtained by replacing I^+_{kl} by I^-_{kl} and q by q^{-1}.

Using the diamond lemma (see, for example, Chapter 4 in [7]), N. Iorgov proved the Poincaré–Birkhoff–Witt theorem for the algebra $U'_q(\mathrm{so}_n)$ (the proof will be published):

Theorem 2. *The elements* ${I^+_{21}}^{m_{21}} {I^+_{31}}^{m_{31}} {I^+_{32}}^{m_{32}} \cdots {I^+_{n1}}^{m_{n1}} \cdots {I^+_{n,n-1}}^{m_{n,n-1}}$, $m_{ij} = 0, 1, 2, \cdots$, *form a basis of the algebra* $U'_q(\mathrm{so}_n)$. *This assertion is true if* I^+_{ij} *are replaced by the corresponding elements* I^-_{ij}.

The algebra $U'_q(\mathrm{so}_n)$ can be embedded into the Drinfeld–Jimbo quantum algebra $U_q(\mathrm{sl}_n)$. This quantum algebra is generated by the elements E_i, F_i, $K_i^{\pm 1} = q^{\pm H_i}$, $i = 1, 2, \cdots, n-1$. Let us introduce the elements

$$\tilde{I}_{j,j-1} = F_{j-1} - q q^{-H_{j-1}} E_{j-1}, \qquad j = 2, 3, \cdots, n.$$

Theorem 3. *The homomorphism* $\varphi : U'_q(\mathrm{so}_n) \to U_q(\mathrm{sl}_n)$ *uniquely determined by the relations* $\varphi(I_{i+1,i}) = \tilde{I}_{i+1,i}$, $i = 1, \cdots, n-1$, *is an isomorphism of* $U'_q(\mathrm{so}_n)$ *to* $U_q(\mathrm{sl}_n)$.

That the mapping $\varphi : U'_q(\mathrm{so}_n) \to U_q(\mathrm{sl}_n)$ is a homomorphism is proved in [4]. In [5] the authors state that it is an isomorphism and say that it can be proved by means of the diamond lemma. However, we could not restore their proof and found another proof. It use the above Poincaré–Birkhoff–Witt theorem for the algebra $U'_q(\mathrm{so}_n)$. Namely, we use the explicit expressions for the elements $I_{ij} \in U'_q(\mathrm{so}_n)$ in terms of the elements of the L-functionals of the quantum algebra $U_q(\mathrm{sl}_n)$ from Subsec. 8.5.2 in [7]. Then we take any linear combination a of the basis elements of $U'_q(\mathrm{so}_n)$ from Theorem 2 and express each element I_{ij} in a in terms of elements of $U_q(\mathrm{sl}_n)$. In this way we obtain the element a expressed in term of elements of $U_q(\mathrm{sl}_n)$. We consider it as a linear combination of basis elements of $U_q(\mathrm{sl}_n)$ given by the Poincaré–Birkhoff–Witt theorem for $U_q(\mathrm{sl}_n)$. Next we show (by using the gradation of $U_q(\mathrm{sl}_n)$ from Subsec. 6.1.5 in [7]) that at least one coefficient in this linear combination is nonvanishing. This proves that φ is an isomorphism from $U'_q(\mathrm{so}_n)$ to $U_q(\mathrm{sl}_n)$.

Corollary. *Let* q *be not a root of unity. Then*

(a) *Finite dimensional irreducible representations of* $U'_q(\mathrm{so}_n)$ *separate elements of this algebra.*

(b) *The homomorphism* $\psi : U'_q(\mathrm{so}_3) \to \hat{U}_q(\mathrm{sl}_2)$ *of the algebra* $U'_q(\mathrm{so}_3)$ *to the extention* $\hat{U}_q(\mathrm{sl}_2)$ *of the quantum algebra* $U_q(\mathrm{sl}_2)$ *(given in [8]) uniquely determined by the relations* $\psi(I_{21}) = \frac{\mathrm{i}}{q-q^{-1}}(q^H - q^{-H})$ *and* $\psi(I_{32}) = (E - F)(q^H + q^{-H})^{-1}$ *is injective.*

The assertion (a) follows from Theorem 3 and from the theorem on separation of elements of the algebra $U_q(\mathrm{sl}_n)$ by its representations (see Subsec. 7.1.5 in [7]). In order to prove (b), we consider representations of $U'_q(\mathrm{so}_3)$ which are obtained by the composition $T \circ \psi$, where T are irreducible finite dimensional representations of $\hat{U}_q(\mathrm{sl}_2)$. It is shown in [8] that every irreducible finite dimensional representation can be obtained from this composition. Now the assertion (b) follows from (a).

Now let us consider finite dimensional irreducible representations of the algebra $U'_q(\mathrm{so}_n)$. If q is not a root of unity, there exists two types of such representations: (a) representations of the classical type (at $q \to 1$ they give the corresponding finite dimensional irreducible representations of the Lie algebra so_n); (b) representations of the nonclassical type (they do not admit the limit $q \to 1$ since in this point the representation operators are singular). These representations are described in [9].

We give irreducible representations of $U'_q(\mathrm{so}_n)$ for q a root of unity. Using a description of central elements of the algebra $U'_q(\mathrm{so}_n)$ for q a root of unity [10] it is proved that every irreducible representation of $U'_q(\mathrm{so}_n)$ in this case is finite dimensional.

Let $q^k = 1$ and k is a smallest positive integer with this property. We fix complex numbers $m_{1,n}, m_{2,n}, ..., m_{\{n/2\},n}$ (here $\{n/2\}$ denotes integral part of $n/2$) and c_{ij}, h_{ij}, $j = 2, 3, \cdots, n-1$, $i = 1, 2, \cdots, \{j/2\}$ such that the differences $h_{ij} - h_{sj}$ and the sums $h_{ij} + h_{sj}$ are not integers. (We also suppose that $h_{p,2p+1}$ are not half-integral.) The set of these numbers will be denoted by ω. Let V be a complex vector space with a basis labelled by the tableaux

$$\{\xi_n\} \equiv \left\{ \begin{matrix} \mathbf{m}_n \\ \mathbf{m}_{n-1} \\ \cdots \\ \mathbf{m}_2 \end{matrix} \right\} \equiv \{\mathbf{m}_n, \xi_{n-1}\} \equiv \{\mathbf{m}_n, \mathbf{m}_{n-1}, \xi_{n-2}\}, \tag{1}$$

where the set of numbers $\mathbf{m}_n$ consists of $\{n/2\}$ numbers $m_{1,n}, m_{2,n}, \cdots, m_{\{n/2\},n}$ given above, and for each $s = 2, 3, \cdots, n-1$, $\mathbf{m}_s$ is a set of numbers $m_{1,s}, \cdots, m_{\{s/2\},s}$ and each $m_{i,s}$ runs independently the values $h_{i,s}, h_{i,s}+1, \cdots, h_{i,s}+k-1$. Thus, $\dim V$ coincides with k^N, where N is the number of positive roots of so_n. It is convenient to use for the numbers $m_{i,s}$ the so-called l-coordinates

$$l_{j,2p+1} = m_{j,2p+1} + p - j + 1, \qquad l_{j,2p} = m_{j,2p} + p - j. \tag{2}$$

To the set of numbers ω there corresponds the irreducible finite dimensional representation T_ω of the algebra $U'_q(\mathrm{so}_n)$. The operator $T_\omega(I_{2p+1,2p})$ of the representation T_ω acts upon the basis elements, labelled by (1), by the formula

$$T_\omega(I_{2p+1,2p})|\xi_n\rangle = \sum_{j=1}^{p} c_{j,2p} \frac{A^j_{2p}(\xi_n)}{q^{l_{j,2p}} + q^{-l_{j,2p}}} \,|(\xi_n)^{+j}_{2p}\rangle - \sum_{j=1}^{p} c^{-1}_{j,2p} \frac{A^j_{2p}((\xi_n)^{-j}_{2p})}{q^{l_{j,2p}} + q^{-l_{j,2p}}} \,|(\xi_n)^{-j}_{2p}\rangle \tag{3}$$

and the operator $T_\omega(I_{2p,2p-1})$ of the representation T_ω acts as

$$T_\omega(I_{2p,2p-1})|\xi_n\rangle = \sum_{j=1}^{p-1} c_{j,2p-1} \frac{B^j_{2p-1}(\xi_n)}{[2l_{j,2p-1}-1][l_{j,2p-1}]} \,|(\xi_n)^{+j}_{2p-1}\rangle$$

$$- \sum_{j=1}^{p-1} c^{-1}_{j,j,2p-1} \frac{B^j_{2p-1}((\xi_n)^{-j}_{2p-1})}{[2l_{j,2p-1}-1][l_{j,2p-1}-1]} \,|(\xi_n)^{-j}_{2p-1}\rangle + \mathrm{i}\, C_{2p-1}(\xi_n)\, |\xi_n\rangle, \tag{4}$$

where numbers in square brackets mean q-numbers: $[a] := (q^a - q^{-a})/(q - q^{-1})$.

In these formulas, $(\xi_n)^{\pm j}_s$ means the tableau (1) in which j-th component $m_{j,s}$ in $\mathbf{m}_s$ is replaced by $m_{j,s} \pm 1$. If $m_{j,s} + 1 = h_{j,s} + k$ (resp. $m_{j,s} - 1 = h_{j,s} - 1$), then we set $m_{j,s}+1=h_{j,s}$ (resp. $m_{j,s}-1=h_{j,s}+k-1$). The coefficients A^j_{2p}, B^j_{2p-1}, C_{2p-1} in (3) and (4) are given by the expressions

$$A^j_{2p}(\xi_n)=\left(\frac{\prod_{i=1}^{p}[l_{i,2p+1}+l_{j,2p}][l_{i,2p+1}-l_{j,2p}-1]\prod_{i=1}^{p-1}[l_{i,2p-1}+l_{j,2p}][l_{i,2p-1}-l_{j,2p}-1]}{\prod_{i\neq j}^{p}[l_{i,2p}+l_{j,2p}][l_{i,2p}-l_{j,2p}][l_{i,2p}+l_{j,2p}+1][l_{i,2p}-l_{j,2p}-1]}\right)^{\frac{1}{2}},$$

$$B^j_{2p-1}(\xi_n)=\left(\frac{\prod_{i=1}^{p}[l_{i,2p}+l_{j,2p-1}][l_{i,2p}-l_{j,2p-1}]\prod_{i=1}^{p-1}[l_{i,2p-2}+l_{j,2p-1}][l_{i,2p-2}-l_{j,2p-1}]}{\prod_{i\neq j}^{p-1}[l_{i,2p-1}+l_{j,2p-1}][l_{i,2p-1}-l_{j,2p-1}][l_{i,2p-1}+l_{j,2p-1}-1][l_{i,2p-1}-l_{j,2p-1}-1]}\right)^{\frac{1}{2}}$$

$$C_{2p-1}(\xi_n)=\frac{\prod_{s=1}^{p}[l_{s,2p}]\prod_{s=1}^{p-1}[l_{s,2p-2}]}{\prod_{s=1}^{p-1}[l_{s,2p-1}][l_{s,2p-1}-1]}.$$

It is seen from (2) that $C_{2p-1}(\xi_n)=0$ if $m_{p,2p}\equiv l_{p,2p}=0$.

The representations T_ω are given by $r=\dim \mathrm{so}_n$ complex numbers. These representations do not exhaust all irreducible representations of $U'_q(\mathrm{so}_n)$ for q a root of unity. The other irreducible representations as well as a proof of the fact that the above formulas determine representations of $U'_q(\mathrm{so}_n)$ will be published.

Acknowledgment

The research of this paper was made possible in part by Award UP1-309 of Civilian Research and Development Foundation and by Award 1.4/206 of Ukrainian DFFD.

References

[1] A. M. Gavrilik and A. U. Klimyk, *Lett. Math. Phys.* **21** (1991), 215.

[2] J. Nelson, T. Regge, and F. Zertuche, *Nucl. Phys.* **B339** (1990), 227.

[3] J. Nelson and T. Regge, *Commun. Math. Phys.* **155** (1993), 561.

[4] M. Noumi, *Adv. Math.* **123** (1993), 16.

[5] M. Noumi, T. Umeda, and M. Wakayama, *Compos. Math.* **104** (1996), 227.

[6] A. M. Gavrilik and N. Z. Iorgov, *Ukr. J. Phys.* **43** (1998), 456.

[7] A. Klimyk and K. Schmüdgen: *Quantum Groups and Their Representations*, Springer, Berlin, 1997.

[8] M. Havlíček, A. Klimyk, and S. Pošta, *J. Math. Phys.* **40** (1999), 2135.

[9] N. Z. Iorgov and A. U. Klimyk, *Czech. J. Phys.* **49** (1999), to be published.

[10] M. Havlíček, A. Klimyk, and S. Pošta, *Czech. J. Phys.* **49** (1999), to be published.